arXiv:1206.6029v1 [astro-ph.CO] 26 Jun 2012

SUPPRESSING HOT GAS ACCRETION TO SUPERMASSIVE BLACK HOLES BY STELLAR WINDS

Shlomi Hillel[1] and Noam Soker[1]

ABSTRACT

We argue that one of the basic assumptions of the Bondi accretion process, that the accreting object has zero pressure, might not hold in many galaxies because of the pressure exerted by stellar winds of star orbiting the central super massive black hole (SMBH). Hence, the Bondi accretion cannot be used in these cases, such as in the galaxy NGC 3115. The winds of these high-velocity stars are shocked to temperatures above the virial temperature of the galaxy, leading to the formation of a hot bubble of size $\sim 0.1 - 10$ pc near the center. This hot bubble can substantially reduce the mass accretion rate by the SMBH. If the density of the hot bubble is lower than that of the interstellar medium (ISM), a density-inversion layer is formed. Adding to other problems of the Bondi process, our results render the Bondi accretion irrelevant for AGN feedback in cooling flow in galaxies and small groups of galaxies and during galaxy formation.

1. INTRODUCTION

It is widely accepted that feedback powered by active galactic nuclei (AGN) has a key role in galaxy formation and in cooling flows in galaxies and in clusters of galaxies. In galaxy formation AGN feedback heats and expels gas (e.g., Ostriker et al. 2010 and references therein), and by that can determine the correlation between the central super-massive black hole (SMBH) mass and some properties of the galaxy. In cooling flow clusters jets launched by the SMBH heat the gas and maintain a small, but non zero cooling flow (see review by McNamara & Nulsen 2007, 2012; Fabian 2012); this is termed a moderate cooling flow.

There is a dispute on how the accretion onto the SMBH occurs, in particular in cooling flows. One camp argues for accretion to be of hot gas via the Bondi accretion process (e.g., Allen et al. 2006; Russell et al. 2010; Narayan & Fabian 2011), while the other side argues that the accretion is of dense and cold clumps in what is termed the cold feedback mechanism (Pizzolato & Soker 2005). The cold feedback mechanism has been strengthened recently by observations of cold gas and by more detailed studies (Revaz et al. 2008; Pope 2009; Wilman et al. 2009, 2011; Nesvadba et al. 2011; Cavagnolo et al. 2011; Gaspari et al. 2012a,b; McCourt et al. 2012; Sharma et al. 2012; Farage et al. 2012; Kashi et al. 2012).

The Bondi accretion process, on the other hand, suffers from two problems. The first problem is that in cooling-flow clusters the Bondi accretion rate is too low to account for the AGN power (e.g., McNamara et al. 2011; Cavagnolo et al. 2011). The second is that there is no time for the feedback to work (Soker et al. 2009). This is because the inflow time scale from even a moderate distance from the center, ~ 1 kpc, in the Bondi accretion process is much longer than the radiative cooling time of the gas. The gas at large distances has no time to communicate with the SMBH before it cools.

In this paper we point out yet another problematic point with the Bondi accretion process. In a recent paper, Wong et al. (2011) resolved the region within the Bondi accretion radius of the S0 galaxy NGC 3115.

---

[1] Department of Physics, Technion – Israel Institute of Technology, Haifa 32000, Israel; shlomihi@tx.technion.ac.il, soker@physics.technion.ac.il

If the density and temperature profile is interpreted as resulting from a Bondi accretion flow onto the $M_{\rm BH} = 2 \times 10^9 M_\odot$ central SMBH, the derived accretion rate is $\dot{M}_B = 2.2 \times 10^{-2} M_\odot~{\rm yr}^{-1}$. They note that for a radiation power of $0.1 \dot{M}_B\, c^2$, the expected accretion luminosity is six orders of magnitude above the observed upper limit. They attribute this to a process where most of the inflowing gas is blown away.

We take a different view. We argue that the Bondi accretion flow is not relevant for the conditions in typical galaxies and clusters of galaxies. The reason is that one cannot assume a zero pressure at the center, either because of stellar winds or because of jets blown by the AGN.

2. THE PRESSURE OF STELLAR WINDS

The pressure exerted by stellar winds of high velocity stars (i.e., moving much faster than the dispersion velocity in the galaxy) with an average mass loss rate per star of $\dot{m}_*$ can be calculated in two limits, which basically lead to the same result. First we calculate the pressure by considering the total outward momentum flux at radius r. The relevant velocity in general is not that of the wind relative to the star, but rather the velocity of the star under the gravitational influence of the SMBH,

$$u_* \simeq \sqrt{\frac{G M_{\rm BH}}{r}} = 2 \times 10^3 \left(\frac{M_{\rm BH}}{10^9 M_\odot} \right)^{1/2} \left(\frac{r}{\rm pc} \right)^{-1/2} ~{\rm km~s}^{-1}. \tag{1}$$

This holds as long as the SMBH gravity dominates that of the galaxy. In NGC 3115 that we study in more detail in section 3, for example, the SMBH gravity dominates that of the galaxy to a distance of ~ 30 pc as the black hole mass is $M_{\rm BH} = 2 \times 10^9 M_\odot$. Let stellar winds from high-velocity stars dominate the pressure inside a sphere of radius R. The pressure exerted by the wind on a surface of radius R is approximately given by adding the ram pressures of winds from all stars inside the sphere of radius R,

$$P_{m*}(R) \simeq n_* \eta \dot{m}_*(R) u_* \frac{4\pi R^3}{3} \frac{1}{4\pi R^2}, \tag{2}$$

where n_* is the stellar density in the center of the galaxy, and η is the fraction of the mass lost by stars that is shocked and heats up. In all our expressions the stellar mass loss rate appears as $\eta \dot{m}_*$.

Some of the mass lost by stars will form dense clumps that will cool rapidly even if being shocked, or will not even be shocked. This is particularly true as most of the mass is being lost by asymptotic giant branch (AGB) stars that have dense winds. The thermal pressure of the ISM in the center will cause part of the winds' gas to form dense clouds. The average mass loss rate is calculated as follows. A solar-like star loses $\sim 0.5 M_\odot$ over $\sim 10^{10}$ yr. Considering an old population of stars, the mass loss rate is lower even. More accurately, most of the mass loss is due to AGB stars, which live for $\sim 10^7$ yr, and lose mass at an average rate of $\sim 10^{-7} M_\odot~{\rm yr}^{-1}$. During the final stages of the AGB the evolution is faster and the mass loss rate is higher. If there is a young stellar population, the total mass loss rate can be much higher. The ram pressure will not increase much beyond few pc because the stellar density decreases.

An alternative point of view would be to express the pressure as (roughly) the energy density of the shocked stellar wind. We also assume a constant pressure and density inside this sphere. This is not the case, but it is adequate for our goals. We can calculate the rate of energy input and multiply by the time it takes the hot gas to leave the inner region

$$\tau_{\rm esc} = \frac{R}{\beta u_*}, \tag{3}$$

where $\beta \lesssim 1$ takes into consideration that the hot gas at the center escapes at velocity lower than the escape velocity. The stellar wind pressure in this case can be written as

$$P_{e*} = \frac{2}{3}\frac{\dot{E}}{V}\frac{R}{\beta u_*}, \tag{4}$$

where the energy deposition rate is

$$\dot{E} = \int_0^R \left(\frac{1}{2} n_* \eta \dot{m}_* u_*^2\right) 4\pi r^2 dr = 2\pi G M_{\rm BH} \eta \dot{m}_* \int_0^R n_*(r) r dr. \tag{5}$$

Scaling the different quantities and assuming a constant stellar density we find

$$P_{e*}(r) = 3 \times 10^{-8} \beta^{-1} \eta \left(\frac{M_{\rm BH}}{10^9 M_\odot}\right)^{1/2} \left(\frac{R}{1 \text{ pc}}\right)^{1/2} \left(\frac{n_*(r)}{5 \times 10^5 \text{ pc}^{-3}}\right) \left(\frac{\dot{m}_*}{10^{-10} M_\odot \text{ yr}^{-1}}\right) \text{ erg cm}^{-3}, \tag{6}$$

where the stellar density is scaled by the average stellar density within ~ 3 pc from the center of NGC 3115 (Kormendy et al. 1996). Equations (4) and (6) are more accurate than equation (2) when the radiative cooling time of the colliding stellar winds is larger than the escape time $\tau_{\rm esc}$, which is the case here due to the high-temperature low-density post-shock stellar winds. Therefore, from now on we will refer to the hot gas region formed by the shocked stellar winds as the hot bubble. For a constant stellar density within radius r, we find $P_{e*} = \frac{3}{2}\beta^{-1} P_{m*}$. If the stellar density drops to zero at some radius r_z (a nonrealistic ideal case), the pressure beyond r_z will drop like $(r/r_z)^{-2}$.

The average density of the hot shocked stellar wind is given by

$$\rho_w \simeq \left(\frac{4\pi}{3} R^3\right)^{-1} \eta \dot{m}_* \frac{R}{\beta u_*(R)} \int_0^R 4\pi n_*(r) r^2 dr \tag{7}$$

The flow structure is schematically drawn in Fig. 1.

3. THE CASE OF NGC 3115

At the Bondi radius $R_B \simeq 210$ pc of the galaxy NGC 3115 the ISM pressure is $P(R_B) = 2 \times 10^{-11}$ erg cm^{-3}, the electron number density is $n_e(R_B) = 0.02$ cm^{-3}, and the temperature is $T(R_B) = 3.5 \times 10^6$ K (Wong et al. 2011). The Bondi radius is given by

$$R_B \simeq \frac{2GM_{\rm BH}}{c_s^2} = 220 \left(\frac{M_{\rm BH}}{2 \times 10^9 M_\odot}\right) \left(\frac{T}{3.5 \times 10^6 \text{ K}}\right) \text{ pc}, \tag{8}$$

where c_s is the sound speed in the undisturbed gas. The temperature and electron density increase inward, reaching values of $T_{20} \simeq 10^7$ K and $n_{e20} \simeq 0.3$ cm^{-3} at $r = 20$ pc (Wong et al. 2011; no values are given at smaller radii). We also note that in NGC 3115 the BH gravity dominates that of the galaxy to a distance of ~ 30 pc as the black hole mass is $M_{\rm BH} = 2 \times 10^9 M_\odot$.

The average density and pressure of the hot bubble according to equations (7) and (6), are drawn in Fig. 2 for a SMBH mass of $M_{\rm BH} = 2 \times 10^9 M_\odot$, and a stellar density given by

$$n_* = 5 \times 10^5 \text{ pc}^{-3} \begin{cases} 1, & r \le 3 \text{ pc} \\ (r/3 \text{ pc})^{-3}, & r > 3 \text{ pc}, \end{cases} \tag{9}$$



Fig. 1.— A schematic drawing (not to scale) of the flow structure where a hot bubble, formed by stellar winds of high-velocity stars orbiting the central SMBH, exerts pressure on the ISM residing outside radius R. If the density in the hot bubble is lower than the ISM density, the flow at R is RT-unstable and a density-inversion layer is formed.

and for $\beta = 1$ (eq. 3) and $\eta = 0.1$ (eq. 2). The density within $r = 3$ pc is from Kormendy et al. (1996), while at $r > 3$ pc is our assumption. The value of the mass loss efficiency, which is the fraction of the mass lost by stars that ends up as hot gas in the hot bubble, is chosen as $\eta = 0.1$ to more or less match the pressure and density of the ISM at $r = 20$ pc. It is a parameter of the model that should be typically in the range of $\sim 0.1 - 1$. The temperature that is calculated from the pressure is also drawn on Fig. 2. Beyond ~ 30 pc the average temperature is only ~ 2 as large as the virial temperature of the cluster, and our assumptions of a hot bubble become inadequate.

The following conclusions emerge from Fig. 2. (1) The pressure of the shocked stellar winds of the high-velocity circum-SMBH stars is larger than the ISM pressure near the center, even for a mass loss efficiency of only $\eta \sim 0.1$. This accounts, we argue, for the accretion rate of NGC 3115 being much lower than the Bondi accretion rate (Wong et al. 2011). (2) At the center, $r < 3$ pc, the rate of mass loss into the hot gas per unit volume is $\dot{\chi} \equiv (n_* \eta \dot{m}_*)_c = 5 \times 10^{-6} M_\odot$ pc^{-3} yr^{-1}. Even if this value is ten times lower, a hot bubble with pressure larger than the ISM pressure of NGC 3115 can still be formed. (3) For $\dot{\chi} \lesssim 10^{-5} M_\odot$ pc^{-3} yr^{-1} the hot bubble's density is lower than that of the ISM. This structure is Rayleigh-Taylor (RT) unstable. This structure is analyzed below.



Fig. 2.— The average density and pressure of the hot bubble according to equations (7) and (6), as well as the temperature that is calculated from the pressure for the stellar density profile given in equation 9. The escape velocity parameter is $\beta = 1$ (eq. 3), and the mass loss parameter of $\eta = 0.1$ (eq. 2) is taken to crudely fit the ISM properties of NGC 3115 at $r = 20$ pc.

4. A TENUOUS HOT BUBBLE FORMED BY STELLAR OR AGN WINDS

We found above that in some cases the hot bubble that formed by the stellar winds of circum-SMBH high-velocity stars can have a lower density than the ISM while its pressure is about equal to the ISM pressure $P_{\rm ISM}$. This situation is prone to RT instability. The same might hold for AGN winds. The power of the winds that is required to form a hot bubble that can support the ISM is

$$W_{\rm wind} \simeq \frac{3}{2} P_{\rm ISM} V \tau_{\rm esc}^{-1} = 3 \times 10^{37} \beta \left(\frac{T n_e}{10^7 \ {\rm K \ cm^{-3}}} \right) \left(\frac{M_{\rm BH}}{10^9 M_\odot} \right)^{1/2} \left(\frac{R}{1 \ {\rm pc}} \right)^{3/2} \ {\rm erg \ s^{-1}}, \qquad (10)$$

where the escape time $\tau_{\rm esc}$ is given by equation (3), and V is the volume of the hot bubble. This implies that even a very weak AGN wind can form such a bubble. With an efficiency of 1%, namely, $W_{\rm wind} = 0.01 \dot{M}_{\rm BH} c^2$, the required accretion rate is $\dot{M}_{\rm BH} = 5 \times 10^{-8} M_\odot \ {\rm yr}^{-1}$. For comparison, we note that the Chandra upper limit on the luminosity of NGC 3115 is $\sim 10^{38}$ erg s^{-1} (Diehl & Statler 2008).

The flow structure considered in this section has the following properties. The hot bubble is continuously supplied by hot gas from the shocked stellar winds or the AGN wind or jets. A pressure equilibrium is maintained between the hot bubble and the ISM, and a structure of a hot tenuous gas supporting a denser and cooler gas is achieved. This structure is RT unstable. Such a structure, we claim, is similar to the density inversion found in the outer atmosphere of red giant stars (e.g., Harpaz 1984; Freytag & Höfner 2008), but not identical. At the outer edge of the recombination zone of hydrogen in red giant stars the convection heat transfer becomes less efficient. The requirement to transfer energy leads to a steep temperature gradient that in turn causes a density inversion, i.e., the density increases outward (e.g., Harpaz 1984). This occurs in the convective region, which is already unstable. In the density-inversion layer in stars, therefore, cold convective cells fall and hot convective cells buoy outward. We suggest that the same process occurs in the flow structure discussed here.

There are some basic differences in the properties of the density-inversion layers of stars and of the case studied here. The main differences are that the hot gas in our case buoys to large distances, and fresh gas from stellar wind or the AGN replaces it. Also, the entire region is optically thin, unlike stars where it is optically thick. In stars the width of the density-inversion region is determined by heat transfer requirements, whereas in our case it is determined by dynamics, mixing, and local heat conduction. In stars the density scale height is not much shorter than the pressure scale height l_p. The size of the convective cells is taken to be of the order of the pressure scale height. In our case the density can change by an order of magnitude from the inner tenuous region to the denser outer ISM, and we expect the RT instability to break the cells to smaller cells. We therefore take the size of the rising and falling gas elements to be $R_c \ll l_p \simeq r$.

We take the density-inversion zone to be of the order of the pressure scale height (in stars it can be much smaller). For a central gravity source the pressure scale height for a constant temperature is given by

$$l_p = r\left[\frac{C_i}{u_*(r)}\right]^2, \tag{11}$$

where C_i is he isothermal sound speed, and u_* is given in equation (1). For an average temperature of the hot gas considered here and the ISM, we can take $l_p \sim r$. Therefore, we assume first that the width of the density-inversion layer is $\Delta r_i \sim r$.

Consider then a spherical parcel of gas (a blob) of radius R_c and density of ρ_b moving with a terminal velocity v_b through an external medium of density ρ_e. The buoyancy force on the blob is

$$F_b = (\rho_e - \rho_c)\frac{4}{3}\pi R_c^3 g. \tag{12}$$

The drag force on the bubble is

$$F_d \approx \frac{1}{2}C_D \pi R_c^2 \rho_e v_t^2, \tag{13}$$

where $C_D \simeq 0.75$ (Kaiser 2003). Assuming $\rho_c \ll \rho_e$ and taking $g = u_*^2/r$, the terminal velocity of the bubble is

$$v_t \approx \left(\frac{8}{3C_D}\right)^{1/2}\left(\frac{R_c}{r}\right)^{1/2} u_* = \beta u_*, \tag{14}$$

where in the second equality we identify the terminal velocity as the velocity by which the hot gas escape from the hot bubble outward, with

$$\beta \simeq 0.6\left(\frac{R_c}{0.1r}\right)^{1/2}. \tag{15}$$

Complex processes take place in the density-inversion layer. (1) Heat conduction time scale over a distance of $\Delta r_T = R_c \sim 0.1$ pc and a temperature difference of $\Delta T = 10^7$ K, is few×10 yr. This is shorter than the fall time of a dense clump from ~ 1 pc. Therefore, the hot bubble gas heats the clump by heat conduction. Closer to the center, the clump will be shredded to smaller cells. Hence, before the dense ISM clumps can reach the center it will be evaporated. This is not true for denser and cooler blobs that fall inward, as in the cold feedback mechanism (Pizzolato & Soker 2005). (2) Because of the stellar motion and/or AGN activity, the density-inversion layer is expected to be more chaotic than just a RT-unstable region. There will be vortices that will increase mixing, namely, reduces the effective value of Δr_T.

5. DISCUSSION AND SUMMARY

We studied the pressure exerted by the winds of circum-SMBH high-velocity stars on the surrounding ISM. We found that in some cases this pressure is significant and can substantially suppress the inflow of the ISM relative to what a simple Bondi accretion would give. Our result can explain the finding of Wong et al. (2011) that the Bondi accretion rate calculated by them from the ISM density and temperature is six orders of magnitude above the observed upper limit on the accretion rate in the S0 galaxy NGC 3115.

In section 3 we quantitatively examined the situation in the galaxy NGC 3115. Shocked winds of circum-SMBH high-velocity stars form a bubble of hot gas whose pressure is significant, as evident from Fig. 2. There are some uncertainties in the model, such as the exact behavior of the stellar mass loss, trajectories of stars around the SMBH, and the stochastic behavior of the post-shock stellar winds. Some of these will be studied in future numerical simulations. However, the result that the stellar winds cannot be ignored is robust. For some values of the parameters we found that a situation might arise where the hot bubble's density is lower than the ISM density. In this case, Rayleigh-Taylor (RT) instability takes place, and a density-inversion layer is formed (see schematic description in Fig. 1). Although hot tenuous gas buoys outward and dense ISM gas moves inward, the density-inversion layer itself continues to exist. The ISM gas is heated near the center and accumulated to the hot bubble.

Our result is more general in showing that in many cases the Bondi accretion process does not work because one of its basic assumptions, that there is no central pressure, breaks down. This is one of several reasons why the Bondi accretion model may not apply in some cases (see section 1).

Finally, we note that our model may be relevant for active galaxies where the hot bubble might be formed by the AGN jets or winds. For typical values of AGN jets and winds the hot bubble density will be low, and a density-inversion layer will be formed. We expect this process to be of high significance in the process of AGN feedback acting in young galaxies. Barring Bondi-like accretion, dense and cold clumps in the ISM can still flow inward and feed the SMBH. Namely, AGN feedback mechanisms require the feeding to be by cold clumps, i.e., a cold feedback mechanism.

This research was supported by the Asher Fund for Space Research and the E. and J. Bishop Research Fund at the Technion, and the Israel Science foundation.

REFERENCES

Allen, S. W., Dunn, R. J. H., Fabian, A. C., Taylor, G. B., & Reynolds, C. S. 2006, MNRAS, 372, 21

Cavagnolo, K. W., McNamara, B. R., Wise, M. W., Nulsen, P. E. J., Brüggen, M., Gitti, M., & Rafferty, D. A. 2011, ApJ, 732, 71

Diehl, S., & Statler, T. S. 2008, ApJ, 680, 897

Fabian, A. C. 2012, ARAA, (arXiv:1204.4114)

Farage, C. L., McGregor, P. J., & Dopita, M. A. 2012, ApJ, 747, 28

Freytag, B., Höfner, S. 2008, A&A, 483, 571

Gaspari, M., Brighenti, F., & Temi, P. 2012a, arXiv:1202.6054

Gaspari, M., Ruszkowski, M., & Sharma, P. 2012b, ApJ, 746, 94

Harpaz, A. 1984, MNRAS, 210, 633

Kaiser, C. R. 2003, MNRAS, 343, 1319

Kashi, A., Nagamine, K., Proga, D., Ostriker, J. P., & Varghese, S. 2012, preprint

Kormendy, J., et al. 1996, ApJ, 459, L57

McCourt, M., Sharma, P., Quataert, E., & Parrish, I. J. 2012, MNRAS, 419, 3319

McNamara, B. R. & Nulsen, P. E. J. 2007, ARA&A, 45, 117

McNamara, B. R., & Nulsen, P. E. J. 2012, New Journal of Physics Focus Issue on Clusters of Galaxies (arXiv:1204.0006)

McNamara, B. R., Rohanizadegan, M., & Nulsen, P. E. J. 2011, ApJ, 727, 39

Narayan, R., & Fabian, A. C. 2011, MNRAS, 926

Nesvadba, N. P. H., Boulanger, F., Lehnert, M. D., Guillard, P., & Salome, P. 2011, A&A, 536, L5

Ostriker, J. P., Choi, E., Ciotti, L., Novak, G. S., & Proga, D. 2010, ApJ, 722, 642

Pizzolato, F. & Soker, N. 2005 ApJ, 632, 821

Pope, E. C. D. 2009, MNRAS, 395, 2317

Revaz, Y., Combes, F., & Salomé, P. 2008, A&A, 477, L33

Russell, H. R., Fabian, A. C., Sanders, J. S., Johnstone, R. M., Blundell, K. M., Brandt, W. N., & Crawford, C. S. 2010, MNRAS, 402, 1561

Sharma, P., McCourt, M., Quataert, E., & Parrish, I. J. 2012, MNRAS, 420, 3174

Soker, N., Sternberg, A., & Pizzolato, F. 2009, in AIP Conf Proc. 1201, The Monsters Fiery Breath: Feedback in Galaxies, Groups & Clusters, ed. S. Heinz & E. Wilcots (Melville, NY: AIP), 321 (arXiv:0909.0220)

Wilman, R. J., Edge, A. C., McGregor, P. J., & McNamara, B. R. 2011, MNRAS, 416, 2060

Wilman, R. J., Edge, A. C., & Swinbank, A. M. 2009, MNRAS, 395, 1355

Wong, K.-W., Irwin, J. A., Yukita, M., Million, E. T., Mathews, W. G., & Bregman, J. N. 2011, ApJ, 736, L23

This preprint was prepared with the AAS LaTeX macros v5.2.